


Malaria 2010



- | | | |
|---|---|---|
| no risk > 2000m | I variable limited risk | II variable elevated risk |
| <ul style="list-style-type: none"> ① JAMMU & KASHMIR ② SIKKIM ③ HIMACHAL PRADESH | <ul style="list-style-type: none"> ④ PUNJAB ⑤ HARYANA ⑥ UTTARANCHAL ⑦ RAJASTHAN ⑧ UTTAR PRADESH ⑨ BIHAR ⑩ ANDHRA PRADESH ⑪ KERLA ⑫ TAMILNADU | <ul style="list-style-type: none"> ⑬ GUJARAT ⑭ MADHYA PRADESH ⑮ CHHATISGARH ⑯ JHARKHAND ⑰ ORISSA ⑱ MAHARASHTRA ⑲ GOA ⑳ KARNATAKA ㉑ BENGAL ㉒ MEGHALAYA ㉓ ARUNACHAL PRADESH ㉔ ASSAM ㉕ NAGALAND ㉖ MANIPUR ㉗ TRIPURA ㉘ MIZORAM ㉙ ANDAMAN & NICOBAR |

(source WHO 2010; UK Malariaguidelines 2007)

Malarone[®] or doxycycline daily; Lariam[®] weekly
 for details : see www.itg.be