

Certificate of additional polio vaccination

Extra measures for twenty-two countries.

13-01-20



Dit is de omschrijving

Polio is an infection caused by a virus. Thanks to vaccination campaigns worldwide, the disease is almost eradicated, except in a few countries.

To avoid the virus spreading from these countries, the World Health Organisation (WHO) has imposed some extra preventive measures.

- All travelers (of all ages) staying more than 4 weeks in countries where WPV1, cVDPV1 or cVPDP3 exist, with risk of international diffusion, have to be vaccinated against polio. This vaccine has to be administered at least 4 weeks and maximum 12 months before leaving one of these countries. Administration of the vaccine has to be documented in the [International Certificate of Vaccination or Prophylaxis](#) (on the pages 4 and 5 reserved for mandatory vaccines):
 - Afghanistan (WPV1)
 - Indonesia (cVDPV1)
 - Malaysia (cVDPV1)
 - Myanmar (cVDPV1)
 - Nigeria (WPV1)
 - Pakistan (WPV1)
 - Philippines (cVDPV1)
- For all travelers staying more than 4 weeks in one of the following countries with risk of cVDPV2, with risk of international diffusion, this supplementary vaccine is recommended but not obliged. This concerns the following countries:
 - Angola
 - Benin
 - Cameroon
 - Central African Republic
 - Chad
 - China
 - DR Congo
 - Ethiopia
 - Ghana
 - Ivory Coast
 - Mozambique
 - Niger
 - Somalia
 - Togo
 - Zambia

Possible vaccines

- Imovax® Polio
- Revaxis®

- Boostrix-Polio®
- Tetravac®
- Infanrix-IPV®
- Hexyon®
- Infanrix-Hexa®

Links

- [International Certificate of Vaccination or Prophylaxis](#)
- [Polioeradication.org](#)
- [Complete WHO report - polio committee](#)